

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
<b>No Management</b>	N	Do not actively manage plants	<p>Minimizing disturbance can protect native species that provide habitat for aquatic fauna; protecting natives may limit spread of invasive species; aquatic plants reduce shoreline erosion and may improve water clarity</p> <p>No immediate financial cost</p> <p>No system disturbance</p> <p>No unintended effects of chemicals</p> <p>Permit not required</p>	<p>May allow small population of invasive plants to become larger, more difficult to control later</p> <p>Excessive plant growth can hamper navigation and recreational lake use</p> <p>May require modification of lake users' behavior and perception</p>
<b>Mechanical Control</b>	May be required under NR 109	Plants reduced by mechanical means	Flexible control	Must be repeated, often more than once per season
		Wide range of techniques, from manual to highly mechanized	Can balance habitat and recreational needs	Can suspend sediments and increase turbidity and nutrient release
a. Handpulling/Manual raking	Y/N	SCUBA divers or snorkelers remove plants by hand or plants are removed with a rake	Little to no damage done to lake or to native plant species	Very labor intensive
		Works best in soft sediments	Can be highly selective	Needs to be carefully monitored
			Can be done by shoreline property owners without permits within an area <30 ft wide OR where selectively removing exotics	Roots, runners, and even fragments of some species, particularly Eurasian watermilfoil (EWM) will start new plants, so all of plant must be removed
			Can be very effective at removing problem plants, particularly following early detection of an invasive exotic species	Small-scale control only

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
b. Harvesting	Y	Plants are "mowed" at depths of 2-5 ft, collected with a conveyor and off-loaded onto shore  Harvest invasives only if invasive is already present throughout the lake	Immediate results  EWM removed before it has the opportunity to autofragment, which may create more fragments than created by harvesting  Minimal impact to lake ecology  Harvested lanes through dense weed beds can increase growth and survival of some fish  Can remove some nutrients from lake	Not selective in species removed  Fragments of vegetation can re-root  Can remove some small fish and reptiles from lake  Initial cost of harvester expensive
<b>Biological Control</b>	Y	Living organisms (e.g. insects or fungi) eat or infect plants	Self-sustaining; organism will over-winter, resume eating its host the next year  Lowers density of problem plant to allow growth of natives	Effectiveness will vary as control agent's population fluctuates  Provides moderate control - complete control unlikely  Control response may be slow  Must have enough control agent to be effective
a. Weevils on EWM	Y	Native weevil prefers EWM to other native water-milfoil	Native to Wisconsin: weevil cannot "escape" and become a problem  Selective control of target species  Longer-term control with limited management	Need to stock large numbers, even if some already present  Need good habitat for overwintering on shore (leaf litter) associated with undeveloped shorelines  Bluegill populations decrease densities through predation

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
b. Pathogens	Y	Fungal/bacterial/viral pathogen introduced to target species to induce mortality	<p>May be species specific</p> <p>May provide long-term control</p> <p>Few dangers to humans or animals</p>	<p>Largely experimental; effectiveness and longevity unknown</p> <p>Possible side effects not understood</p>
c. Allelopathy	Y	Aquatic plants release chemical compounds that inhibit other plants from growing	<p>May provide long-term, maintenance-free control</p> <p>Spikerushes (<i>Eleocharis</i> spp.) appear to inhibit Eurasian watermilfoil growth</p>	<p>Initial transplanting slow and labor-intensive</p> <p>Spikerushes native to WI, and have not effectively limited EWM growth</p> <p>Wave action along shore makes it difficult to establish plants; plants will not grow in deep or turbid water</p>
d. Planting native plants	Y	Diverse native plant community established to repel invasive species	<p>Native plants provide food and habitat for aquatic fauna</p> <p>Diverse native community may be "resistant" to invasive species</p> <p>Supplements removal techniques</p>	<p>Initial transplanting slow and labor-intensive</p> <p>Nuisance invasive plants may outcompete plantings</p> <p>Largely experimental; few well-documented cases</p> <p>If transplants from external sources (another lake or nursery), may include additional invasive species or "hitchhikers"</p>

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
<b>Physical Control</b>	Required under Ch. 30 / NR 107	Plants are reduced by altering variables that affect growth, such as water depth or light levels		
a. Fabrics/ Bottom Barriers	Y	Prevents light from getting to lake bottom	<p>Reduces turbidity in soft-substrate areas</p> <p>Useful for small areas</p>	<p>Eliminates all plants, including native plants important for a healthy lake ecosystem</p> <p>May inhibit spawning by some fish</p> <p>Need maintenance or will become covered in sediment and ineffective</p> <p>Gas accumulation under blankets can cause them to dislodge from the bottom</p> <p>Affects benthic invertebrates</p> <p>Anaerobic environment forms that can release excessive nutrients from sediment</p>
b. Drawdown	Y, May require Environmental Assessment	<p>Lake water lowered with siphon or water level control device; plants killed when sediment dries, compacts or freezes</p> <p>Season or duration of drawdown can change effects</p>	<p>Winter drawdown can be effective at restoration, provided drying and freezing occur. Sediment compaction is possible over winter</p> <p>Summer drawdown can restore large portions of shoreline and shallow areas as well as provide sediment compaction</p> <p>Emergent plant species often rebound near shore providing fish and wildlife habitat, sediment stabilization, and increased water quality</p> <p>Success demonstrated for reducing EWM, variable success for curly-leaf pondweed (CLP)</p> <p>Restores natural water fluctuation important for all aquatic ecosystems</p>	<p>Plants with large seed bank or propagules that survive drawdown may become more abundant upon refilling</p> <p>May impact attached wetlands and shallow wells near shore</p> <p>Species growing in deep water (e.g. EWM) that survive may increase, particularly if desirable native species are reduced</p> <p>Can affect fish, particularly in shallow lakes if oxygen levels drop or if water levels are not restored before spring spawning</p> <p>Winter drawdown must start in early fall or will kill hibernating reptiles and amphibians</p> <p>Navigation and use of lake is limited during drawdown</p>

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
c. Dredging	Y	<p>Plants are removed along with sediment</p> <p>Most effective when soft sediments overlay harder substrate</p> <p>For extremely impacted systems</p> <p>Extensive planning required</p>	<p>Increases water depth</p> <p>Removes nutrient rich sediments</p> <p>Removes soft bottom sediments that may have high oxygen demand</p>	<p>Severe impact on lake ecosystem</p> <p>Increases turbidity and releases nutrients</p> <p>Exposed sediments may be recolonized by invasive species</p> <p>Sediment testing may be necessary</p> <p>Removes benthic organisms</p> <p>Dredged materials must be disposed of</p>
d. Dyes	Y	<p>Colors water, reducing light and reducing plant and algal growth</p>	<p>Impairs plant growth without increasing turbidity</p> <p>Usually non-toxic, degrades naturally over a few weeks</p>	<p>Appropriate for very small water bodies</p> <p>Should not be used in pond or lake with outflow</p> <p>Impairs aesthetics</p> <p>Effects to microscopic organisms unknown</p>
e. Non-point source nutrient control	N	<p>Runoff of nutrients from the watershed are reduced (e.g. by controlling construction erosion or reducing fertilizer use) thereby providing fewer nutrients available for plant growth</p>	<p>Attempts to correct source of problem, not treat symptoms</p> <p>Could improve water clarity and reduce occurrences of algal blooms</p> <p>Native plants may be able to better compete with invasive species in low-nutrient conditions</p>	<p>Results can take years to be evident due to internal recycling of already-present lake nutrients</p> <p>Requires landowner cooperation and regulation</p> <p>Improved water clarity may increase plant growth</p>

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
<b>Chemical Control</b>	Y, Required under NR 107	<p>Granules or liquid chemicals kill plants or cease plant growth; some chemicals used primarily for algae</p> <p>Results usually within 10 days of treatment, but repeat treatments usually needed</p> <p>Chemicals must be used in accordance with label guidelines and restrictions</p>	<p>Some flexibility for different situations</p> <p>Some can be selective if applied correctly</p> <p>Can be used for restoration activities</p>	<p>Possible toxicity to aquatic animals or humans, especially applicators</p> <p>May kill desirable plant species, e.g. native water-milfoil or native pondweeds; maintaining healthy native plants important for lake ecology and minimizing spread of invasives</p> <p>Treatment set-back requirements from potable water sources and/or drinking water use restrictions after application, usually based on concentration</p> <p>May cause severe drop in dissolved oxygen causing fish kill, depends on plant biomass killed, temperatures and lake size and shape</p> <p>Often controversial</p>
a. 2,4-D	Y	<p>Systemic<sup>1</sup> herbicide selective to broadleaf<sup>2</sup> plants that inhibits cell division in new tissue</p> <p>Applied as liquid or granules during early growth phase</p>	<p>Moderately to highly effective, especially on EWM</p> <p>Monocots, such as pondweeds (e.g. CLP) and many other native species not affected</p> <p>Can be selective depending on concentration and seasonal timing</p> <p>Can be used in synergy with endothall for early season CLP and EWM treatments</p> <p>Widely used aquatic herbicide</p>	<p>May cause oxygen depletion after plants die and decompose</p> <p>May kill native dicots such as pond lilies and other submerged species (e.g. coontail)</p> <p>Cannot be used in combination with copper herbicides (used for algae)</p> <p>Toxic to fish</p>

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
b. Endothall	Y	Broad-spectrum <sup>3</sup> , contact <sup>4</sup> herbicide that inhibits protein synthesis  Applied as liquid or granules	Especially effective on CLP and also effective on EWM  May be effective in reducing reestablishment of CLP if reapplied several years in a row in early spring  Can be selective depending on concentration and seasonal timing  Can be combined with 2,4-D for early season CLP and EWM treatments, or with copper compounds  Limited off-site drift	Kills many native pondweeds  Not as effective in dense plant beds; heavy vegetation requires multiple treatments  Not to be used in water supplies; post-treatment restriction on irrigation  Toxic to aquatic fauna (to varying degrees)
c. Diquat	Y	Broad-spectrum, contact herbicide that disrupts cellular functioning  Applied as liquid, can be combined with copper treatment	Mostly used for water-milfoil and duckweed  Rapid action  Limited direct toxicity on fish and other animals	May impact non-target plants, especially native pondweeds, coontail, elodea, naiads  Toxic to aquatic invertebrates  Must be reapplied several years in a row  Ineffective in muddy or cold water (<50°F)
d. Fluridone	Y; special permit and Environmental Assessment may be required	Broad-spectrum, systemic herbicide that inhibits photosynthesis  Must be applied during early growth stage  Available with a special permit only; chemical applications beyond 150 ft from shore not allowed under NR 107  Applied at very low concentration at whole lake scale	Effective on EWM for 1 to 4 years with aggressive follow-up treatments  Some reduction in non-target effects can be achieved by lowering dosage  Slow decomposition of plants may limit decreases in dissolved oxygen  Low toxicity to aquatic animals	Affects non-target plants, particularly native milfoils, coontails, elodea, and naiads, even at low concentrations  Requires long contact time at low doses: 60-90 days  Demonstrated herbicide resistance in hydrilla subjected to repeat treatments  In shallow eutrophic systems, may result in decreased water clarity  Unknown effect of repeat whole-lake treatments on lake ecology

# Management Options for Aquatic Plants



Draft updated Oct 2006

Option	Permit Needed?	How it Works	PROS	CONS
e. Glyphosate	Y	Broad-spectrum, systemic herbicide that disrupts enzyme formation and function  Usually used for purple loosestrife stems or cattails  Applied as liquid spray or painted on loosestrife stems	Effective on floating and emergent plants such as purple loosestrife  Selective if carefully applied to individual plants  Non-toxic to most aquatic animals at recommended dosages  Effective control for 1-5 years	RoundUp is often incorrectly substituted for Rodeo - Associated surfactants of RoundUp believed to be toxic to reptiles and amphibians  Cannot be used near potable water intakes  Ineffective in muddy water  No control of submerged plants
f. Triclopyr	Y	Systemic herbicide selective to broadleaf plants that disrupts enzyme function  Applied as liquid spray or liquid	Effective on many emergent and floating plants  More effective on dicots, such as purple loosestrife; may be more effective than glyphosate  Control of target plants occurs in 3-5 weeks  Low toxicity to aquatic animals  No recreational use restrictions following treatment	Impacts may occur to some native plants at higher doses (e.g. coontail)  May be toxic to sensitive invertebrates at higher concentrations  Retreatment opportunities may be limited due to maximum seasonal rate (2.5 ppm)  Sensitive to UV light; sunlight can break herbicide down prematurely  Relatively new management option for aquatic plants (since 2003)
g. Copper compounds	Y	Broad-spectrum, systemic herbicide that prevents photosynthesis  Used to control planktonic and filamentous algae  Wisconsin allows small-scale control only	Reduces algal growth and increases water clarity  No recreational or agricultural restrictions on water use following treatment  Herbicidal action on hydrilla, an invasive plant not yet present in Wisconsin	Elemental copper accumulates and persists in sediments  Short-term results  Long-term effects of repeat treatments to benthic organisms unknown  Toxic to invertebrates, trout and other fish, depending on the hardness of the water  Clear water may increase plant growth

<sup>1</sup>Systemic herbicide - Must be absorbed by the plant and moved to the site of action. Often slower-acting than contact herbicides.  
<sup>2</sup>Broadleaf herbicide - Affects only dicots, one of two groups of plants. Aquatic dicots include waterlilies, bladderworts, watermilfoils, and coontails.  
<sup>3</sup>Broad-spectrum herbicide - Affects both monocots and dicots.  
<sup>4</sup>Contact herbicide - Unable to move within the plant; kills only plant tissue it contacts directly.  
 Specific effects of herbicide treatments dependent on timing, dosage, duration of treatment, and location.  
 References to registered products are for your convenience and not intended as an endorsement or criticism of that product versus other similar products.  
**This document is intended to be a guide to available aquatic plant control techniques, and is not necessarily an exhaustive list.**  
**Please contact your local Aquatic Plant Management Specialist when considering a permit.**